



Deglaciation chronology of the Donegal Ice Centre, northwest Ireland.

Wilson, P., Ballantyne, C. K., Benetti, S., Small, D., Fabel, D., & Clark, C. D. (2019). Deglaciation chronology of the Donegal Ice Centre, northwest Ireland. *Journal of Quaternary Science*, 34(1), 16-28.
<https://doi.org/10.1002/jqs.3077>

[Link to publication record in Ulster University Research Portal](#)

Published in:
Journal of Quaternary Science

Publication Status:
Published (in print/issue): 11/01/2019

DOI:
[10.1002/jqs.3077](https://doi.org/10.1002/jqs.3077)

Document Version
Author Accepted version

General rights
Copyright for the publications made accessible via Ulster University's Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy
The Research Portal is Ulster University's institutional repository that provides access to Ulster's research outputs. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact pure-support@ulster.ac.uk.

Journal of Quaternary Science

Deglaciation chronology of the Donegal Ice Centre, northwest Ireland.

Journal:	<i>Journal of Quaternary Science</i>
Manuscript ID	JQS-18-0070
Wiley - Manuscript type:	Research Article
Date Submitted by the Author:	25-Jun-2018
Complete List of Authors:	Wilson, Peter; University of Ulster, School of Environmental Sciences Ballantyne, Colin; University of St Andrews, Geography and Geosciences Benetti, Sara; University of Ulster, School of Environmental Sciences Small, David; Durham University, Department of Geography Fabel, Derek; University of Glasgow, Geographical and Earth Sciences Clark, Chris; Univ Sheffield, Dept Geography
Keywords:	British-Irish Ice Sheet, deglaciation, terrestrial cosmogenic nuclide surface exposure dating, northwest Ireland, isochrones

SCHOLARONE™
Manuscripts

**Deglaciation chronology of the Donegal Ice Centre,
northwest Ireland**

**PETER WILSON¹, COLIN K. BALLANTYNE², SARA BENETTI¹, DAVID SMALL³,
DEREK FABEL⁴, CHRIS D. CLARK⁵**

¹School of Geography and Environmental Sciences, Ulster University, Coleraine BT52 1SA, UK.

²School of Geography and Sustainable Development, University of St. Andrews, St Andrews
KY16 9AL, UK.

³Department of Geography, University of Durham, Durham DH UK.

⁴Scottish Universities Environmental Research Centre, East Kilbride G75 0QF, UK

⁵Department of Geography, University of Sheffield, Sheffield S10 2TN, UK

Corresponding author: Peter Wilson, as above.
Email address: P.Wilson@ulster.ac.uk

ABSTRACT:

During the Last Glacial Maximum, Donegal in NW Ireland functioned as an independent centre of ice dispersal that separated and fed into the Donegal Bay Ice Stream (sourced in the Irish Midlands) to the south and the Hebrides/Malin Sea Ice Stream to the north. We report geochronological data that demonstrate marked contrasts in the timing and rate of deglaciation in northern and southern Donegal. In northern Donegal, which occupied an inter-ice-stream location, decoupling from the Hebrides/Malin Sea Ice Stream resulted in formation of a marine embayment along the north coast by $\sim 22\text{--}21$ ka, and subsequent slow ($\sim 4 \pm 1$ m a $^{-1}$) climatically-driven inland retreat of the ice margin to mountain source areas by ~ 17 ka. By contrast in southern Donegal, which lay near the axis of the Donegal Bay Ice Stream, deglaciation was delayed until ~ 18 ka following readvance of ice to a moraine in outer Donegal Bay. The ice margin subsequently underwent net retreat, apparently uninterrupted by readvances, at a net rate of $\sim 18 \pm 6$ m a $^{-1}$. A mean terrestrial cosmogenic nuclide age of ~ 14.8 ka obtained for samples from the foothills of the Blue Stack Mountains in SE Donegal indicates that ice persisted in valley heads and cirques at the beginning of the Lateglacial Interstadial, suggesting that these and nearby mountains supported the last remnants of the Irish Ice Sheet prior to complete deglaciation of Ireland, and that almost all of the shrinkage of the ice sheet in this sector occurred under stadial conditions prior to the onset of interstadial warming at ~ 14.7 ka.

KEYWORDS: British-Irish Ice Sheet, deglaciation, terrestrial cosmogenic nuclide surface exposure dating, northwest Ireland

Introduction

During the last (Late Devensian / Late Midlandian) ice-sheet glaciation of Britain and Ireland, (~32–15 ka) the mountains of Donegal in northwest Ireland formed an independent centre of ice dispersal within the more extensive British-Irish Ice Sheet (BIIS). Ice radiating from the Donegal Ice Centre fed north and northwest into a major ice stream (the Hebrides/Malin Sea Ice Stream) on the adjacent Malin Shelf, west and southwest into Donegal Bay, and to the east was confluent with the ice occupying the Irish Midlands. The Donegal ice dome was therefore pivotal in separating ice flows from the Irish Midlands and western Scotland (Fig. 1). Radial ice flow over Donegal for at least part of the last glaciation is demonstrated by the distribution of local erratics, absence of allochthonous erratics, and the alignments of drumlins, moraines, roches moutonnées, striae and meltwater channels (Charlesworth, 1924; Dury, 1957, 1958, 1964; Stephens and Synge, 1965; Colhoun, 1973; McCabe *et al.*, 1993; Knight and McCabe, 1997; Ballantyne *et al.*, 2007; Smith and Knight, 2011; Knight, 2012). Flowsets reconstructed by Greenwood and Clark (2009b) show that ice moving south from the Donegal Ice Centre was confluent with west-flowing ice from the Irish Midlands in Donegal Bay, forming an ice stream (the Donegal Bay Ice Stream) that extended northwestwards towards the shelf edge.

Geophysical data obtained for the adjacent offshore shelves indicate that at the global Last Glacial Maximum (gLGM; 26.5-19 ka, P.U. Clark *et al.*, 2009)) grounded ice extended as far as the shelf break, ~100 km to the west, where it terminated in a marine setting (Benetti *et al.*, 2010; Dunlop *et al.* 2010; Ó Cofaigh *et al.*, 2012). These data also indicate that Donegal ice coalesced with ice from western Scotland ~60 km north of the present Donegal coastline. Donegal is therefore the key location for determining the timing of decoupling of Irish- and Scottish-sourced

ice during the last deglaciation, and is also important for establishing the chronology of ice retreat after the ice margin had retreated to the present coastline.

As part of the wider BRITICE-CHRONO project (<http://www.sheffield.ac.uk/geography/research/britice-chrono/home>), designed to establish a detailed deglaciation chronology of the last BIIS, we present 20 new ^{10}Be and ^{36}Cl terrestrial cosmogenic nuclide (TCN) surface exposure ages from six sites in Donegal that were selected to complement and extend the existing deglaciation chronology. The aims of this paper are: (1) to establish the timing of the decoupling of Scottish and Irish ice flowing west across the Malin Shelf; (2) to reconstruct the chronology of ice margin retreat in Donegal Bay; (3) to determine the net rate of ice-margin recession inland from the north coast of Donegal and in Donegal Bay; (4) to establish for how long ice persisted locally in the Donegal mountains following its retreat from coastal lowlands; and (5) to explore the wider implications of our results for the interpretation of the deglaciation chronology of the western sector of the last BIIS. The chronology of offshore ice margin retreat from the shelf edge towards the present coastline is considered in a separate paper based on new radiocarbon ages obtained from marine microfauna retrieved from sediment cores along a transect from the shelf edge to the outer part of Donegal Bay (Ó Cofaigh *et al.*, 2018).

Donegal

Regional setting and ice dome extent

County Donegal (54°28'–55°22' N, 06°55'–08°46' W) is predominantly underlain by granites, quartzites and schists with a pronounced northeast-southwest structural grain that has been accentuated by repeated episodes of Quaternary glacial erosion (Long and McConnell, 1997, 1999). The north and west of the county are mountainous; many summits exceed 500 m OD with

the highest point (Errigal) at 751 m OD. From detailed mapping of erosional and depositional landforms, Charlesworth (1924) proposed that the Donegal mountains had nourished and maintained an independent ice dome during the last glaciation, and placed the former ice divide along a line running approximately north-south from the Derryveagh Mountains to the Blue Stack Mountains, close to the present watershed (Fig. 1). He showed that ice-flow from this elongated dome was essentially radial with a focus along pre-existing structurally-controlled valleys, but argued that during maximum ice extent topography was probably less of a constraining influence on ice-flow directions than during build-up and retreat phases. Subsequent work is generally supportive of this ice dome hypothesis (e.g. Ballantyne *et al.*, 2007; Greenwood and Clark, 2009; Smith and Knight, 2011).

In contrast, the thickness attained by the ice dome has been a contentious issue. Charlesworth (1924) claimed that all summits lay beneath the ice, although it is not clear if he was referring to the local Last Glacial Maximum (ILGM), which is placed at ~26.3–24.8 ka at the shelf break to the west of Donegal (Ó Cofaigh *et al.*, 2018). Complete burial of the mountains by the last ice sheet was favoured by McCabe (1995), while Sellier (1995) maintained that areas above ~550 m OD in the Derryveagh Mountains had remained ice free. On the basis of geomorphological evidence and clay-fraction mineralogy, Ballantyne *et al.* (2007) argued for an ice-shed altitude in excess of 700 m OD, but also reported an absence of evidence for glacial modification on six peripheral summits, including Errigal, and regarded these as being either nunataks during the ILGM or buried beneath a cover of non-erosive cold-based ice.

Conflicting interpretations also concern the lateral extent of the Donegal ice dome. Although Charlesworth (1924) envisaged ice extending offshore to the north and west, others have placed the limit onshore in the north of the county (Stephens and Synge, 1965; Bowen *et al.*, 2002), and a limited offshore extent of ~10-30 km to the west has been suggested (e.g. McCabe, 1985;

Bowen *et al.*, 1986; Knight, 2003; Ballantyne *et al.*, 2007). More recent work utilising geophysical techniques to image seabed topography has demonstrated that a concentric sequence of nested moraine ridges indicative of deposition by a grounded ice mass extends westwards to the shelf break 90–100 km from the west coast of Donegal (Sejrup *et al.*, 2005; Benetti *et al.*, 2010; Dunlop *et al.* 2010; Ó Cofaigh *et al.*, 2012). Radiocarbon dates obtained for marine microfauna in cores retrieved from sediments on the Atlantic shelf northwest of Ireland confirm that these moraines were deposited at the margin of the last ice sheet, and indicate that ice nourished in Donegal began to retreat from the shelf edge in the interval between 26.3 and 24.8 ka cal BP (Ó Cofaigh *et al.* 2018).

Recognition that the last ice sheet extended to the edge of the Malin Shelf strongly suggests that the Donegal ice dome was of sufficient thickness to have buried all mountain summits during the ILGM, a proposition also supported by climate-proxy-driven thermo-mechanical models of ice-sheet build-up and decay (Hubbard *et al.*, 2009). Support for this premise comes from southwest Ireland where Ballantyne *et al.* (2011) have argued that the Kerry-Cork Ice Cap attained an altitude of at least 1200 m OD, >200 m above the highest summits, and from northwest Scotland where Fabel *et al.* (2012) have demonstrated that the last ice sheet overtopped all mountain summits. It is therefore extremely unlikely that any of the mountain summits in Donegal formed palaeonunataks during the ILGM (Ballantyne and Ó Cofaigh, 2017).

Legacy ages, related BRITICE-CHRONO ages and deglaciation

Several previous studies have utilised either TCN (cosmogenic ^{10}Be or ^{36}Cl) surface exposure dating or ^{14}C dating to establish the timing of ice retreat and/or readvance from sites in Donegal. Collectively these ages provide the foundation of a deglaciation chronology (Fig.1; Table 1).

1
2
3 148 Cosmogenic ^{10}Be exposure ages cited here have been recalibrated using the Loch Lomond
4
5 149 Production Rate (LLPR), and are followed in brackets by the equivalent ages obtained from the
6
7
8 150 CRONUScale online calculator; details of these procedures are given in the next section. The ^{14}C
9
10 151 ages have been (re)calibrated using OxCal 4.2 and, for marine-derived samples, the Marine-13
11
12 152 dataset with a marine reservoir correction of 400 years (Bronk Ramsey, 2009; Reimer *et al.*, 2013).
13
14 153 All ^{14}C ages are reported to two decimal places as cal ka BP; TCN ages are reported to one decimal
15
16
17 154 place as ka. Mean ages reported for two or more TCN ages below and in Tables 1 and 3 are
18
19 155 uncertainty-weighted means.

20
21 156 Bowen *et al.* (2002) obtained ^{36}Cl ages of 25.1 ± 1.1 ka, from glacially-smoothed quartzite
22
23 157 bedrock at Malin Head, and 31.0 ± 17.0 ka, from either a glacially-transported granite boulder or
24
25
26 158 bedrock at Bloody Foreland, but the large uncertainty on the latter age prevents meaningful
27
28 159 interpretation, and the former is probably compromised by nuclide inheritance (Ballantyne and Ó
29
30
31 160 Cofaigh, 2017). For Corvish, at the head of Trawbreaga Bay on the north coast, McCabe and Clark
32
33 161 (2003) reported ^{14}C ages for marine microfaunas within *in-situ* and deformed marine sediments.
34
35 162 The basal *in-situ* laminated muds yielded ages of 20.68 ± 0.16 cal ka BP and 18.24 ± 0.13 cal ka BP;
36
37
38 163 the older date implies initial deglaciation before ~ 20.7 cal ka BP. Overlying deformed sands and
39
40 164 muds gave ages of 19.50 ± 0.50 , 18.32 ± 0.18 and 19.03 ± 0.19 cal ka BP, and were interpreted by
41
42 165 McCabe and Clark (2003) as evidence for reworking of the underlying laminated muds by ice
43
44
45 166 readvance at ~ 18 ka. An age of 17.06 ± 0.18 cal. ka BP from *in-situ* rhythmically bedded marine
46
47 167 muds overlying the deformed muds was regarded as minimal for final deglaciation of the bay.

48
49 168 Seven consistent ^{10}Be exposure ages from glacially-transported boulders on a lateral
50
51 169 moraine at Bloody Foreland, the northwesternmost point of Donegal, have given an uncertainty-
52
53
54 170 weighted mean age of 21.8 ± 1.2 ka (21.1 ± 1.8 ka) (Ballantyne *et al.*, 2007; Clark *et al.*, 2009a;
55
56 171 Ballantyne and Ó Cofaigh, 2017; Fig. 2a). Two ^{10}Be exposure ages from bedrock and a glacially-

transported boulder on Aran Island, 20 km southwest of Bloody Foreland, have yielded an uncertainty-weighted mean age of 21.7 ± 1.1 ka (21.0 ± 1.7 ka) (Cullen, 2013). The consistency of these two mean ages provides strong support for retreat of the ice-sheet margin across the Irish sector of the Malin Shelf between ~ 26 – 25 ka and ~ 22 – 21 ka (Clark *et al.*, 2012; Ó Cofaigh *et al.*, 2012, 2018), decoupling of Malin Shelf ice from Donegal Bay ice at ~ 22 – 21 ka and the beginning of ice retreat at that time from the present coast towards the mountains. At Glencolumbkille in southwest Donegal, ^{10}Be exposure ages of 17.9 ± 0.9 ka (17.4 ± 1.4 ka) and 19.8 ± 1.0 ka (19.1 ± 1.5 ka) from vein quartz in, respectively, a glacially-transported boulder and a roche moutonnée were reported by Ballantyne *et al.* (2007). The latter age overlaps within 1σ uncertainties with the mean values from Bloody Foreland and Aran Island, but may be compromised by nuclide inheritance (see below).

The timing of deglaciation of the mountains of Donegal is indicated by ^{10}Be exposure ages for two sites. Glacially-plucked bedrock at 405–430 m OD on a col to the east of Errigal in north Donegal has produced three consistent ^{10}Be exposure ages averaging 17.8 ± 0.9 ka (17.6 ± 1.4 ka) and a minimum age for deglaciation of Slieve League in southwest Donegal is provided by three consistent ^{10}Be exposure ages averaging 17.1 ± 0.9 ka (16.9 ± 1.4 ka) obtained for samples from rockslide runout debris (Ballantyne *et al.*, 2013b).

A ^{14}C age of 15.38 ± 0.12 cal ka BP from the basal organic mud of Lough Nadourcan (Watson *et al.*, 2010) provides a minimum age for deglaciation of the low ground along the eastern margin of the Derryveagh Mountains. However, this age is ~ 700 years earlier than the rapid warming identified in the Greenland ice core records and INTIMATE event stratigraphy as marking the onset of the Lateglacial Interstadial at ~ 14.7 ka (Rasmussen *et al.*, 2014), suggesting that the Lough Nadourcan basal ^{14}C age may be compromised by the incorporation of reworked

older carbon. Nevertheless, it is unlikely that ice on low ground survived much beyond the start of interstadial warming even if small glaciers persisted in the mountains.

Legacy ages from sites in north Mayo, along the south side of Donegal Bay, and BRITICE-CHRONO ages from Donegal Bay (Fig. 1) are relevant to the deglaciation chronology of south Donegal, and therefore are also considered here. McCabe *et al.* (1986, 2005) reported eight ¹⁴C ages obtained for marine shells and foraminifera within glacimarine sediments at Fiddauntawnanoneen and Belderg Pier on the north coast of Mayo. Seven of these ages range between 20.38±0.31 cal ka BP and 19.16±0.21 cal ka BP; the remaining age (22.09±0.28 cal. ka BP) is significantly older and may indicate the reworking of older sediment (J. Clark *et al.*, 2012). Deglaciation of these adjacent sites and, by inference, the outer reaches of Donegal Bay, therefore appears to have occurred around or slightly before ~20 ka (Ballantyne and Ó Cofaigh, 2017).

To the northeast of these two sites, a distinct ice margin position is represented by the Donegal Bay Moraine (DBM), an offshore moraine that extends for 35 km north-south across outer Donegal Bay (Benetti *et al.*, 2010; Ó Cofaigh *et al.* 2012). Deformation of stratified glacimarine deposits indicates that the moraine represents a readvance of the ice margin. Radiocarbon ages for mixed benthic foraminifera within glacimarine sediments in 76–99 m water depth on either side of the moraine (Fig. 1) constrain moraine formation to between 20.24±0.24 cal ka BP and 17.92±0.16 cal ka BP (Ó Cofaigh *et al.*, 2018), and moraine formation at 20–19 ka was inferred by Ó Cofaigh *et al.* (2018).

Finally, eight cosmogenic ¹⁰Be exposure ages from vein quartz in glacially-transported boulders at three sites associated with the Tawnywaddyduff moraine system on the northern slopes of the Ox Mountains south of Sligo Bay (Fig. 1) returned ages ranging from 21.1±1.7 ka (21.0±2.3 ka) to 15.7±1.6 ka (16.0±2.0 ka). The overall average of these ages (~18 ka) was taken by Clark *et al.* (2009b) to represent the timing of a readvance of the ice sheet and construction of the moraine.

However, Ballantyne and Ó Cofaigh (2017) questioned this conclusion, noting that the age range spanned >5 ka and that two distinct age groupings are represented, with three older ages (mean 20.3 ± 1.3 ka (20.3 ± 1.9 ka)) and five younger ages (mean 16.6 ± 1.0 ka (16.7 ± 1.5 ka)). The older sample ages are from a site on the west side of the Ox Mountains; four of the younger ages are from the east side and the other age came from the northern slopes.

Field sites and methods

The six sites sampled for TCN surface exposure dating were selected to provide deglaciation-age transects along the north and south coasts of Donegal and an additional deglaciation age for the northern (Derryveagh) mountains (Fig. 1). For the north coast transect we sampled on the headlands of Rosguill and Malin Head, respectively 29 km and 63 km northeast of the Bloody Foreland site dated by Ballantyne *et al.* (2007) and Clark *et al.* (2009a). For the southern transect we sampled at Glencolumbkille, close to the western extremity of the Slieve League peninsula, at Kilcar on the SW coast of Donegal, and on the lower southern slopes of the Blue Stack Mountains. The latter two sites are, respectively, 14 km southeast and 41 km east of Glencolumbkille. In the northern mountains we obtained samples from a prominent valley-floor boulder limit in the Poisoned Glen, Derryveagh Mountains.

Samples were collected from the upper surface of large, glacially-deposited boulders or ice-scoured bedrock using a hammer and chisel. Twelve boulder samples comprised whole rock (granite, conglomerate sandstone or dolerite), four were from protruding quartz veins in quartzite or schist boulders, two consisted of quartz pebbles embedded in conglomerate boulders, and two samples were from quartzite bedrock (Fig. 2, Table 2). A compass and clinometer were used to record the geometry of the sampled surfaces and the skyline topography. Locations and altitudes

were determined with a hand-held GPS unit cross-referenced to a 1:50,000 topographic map. Sample thickness was measured using callipers, density was determined by the displacement of sub-samples in water, and topographic shielding was calculated using the CRONUS-Earth online calculator (Table 2).

Samples were processed for cosmogenic ^{10}Be and ^{36}Cl analysis at the NERC Cosmogenic Isotope Analysis Facility (CIAF). For ^{10}Be , samples were crushed and sieved to 250-500 μm and quartz was separated in a Frantz® isodynamic magnetic mineral separator. before being repeatedly etched with HF (Kohl and Nishiizumi, 1992). Purified quartz was spiked with ~ 0.2 mg of ^9Be and dissolved. Be was extracted and isolated following the methodology described in Child *et al.* (2000) before being, precipitated as $\text{Be}(\text{OH})_2$ and baked to BeO in a quartz crucible. BeO was mixed with Nb and pressed into a copper cathode. For ^{36}Cl , samples were crushed and sieved to <500 μm , leached in hot HNO_3 (trace metal analysis grade) and then washed thoroughly with ultrapure water to remove meteoric ^{36}Cl contamination from grain surfaces. Each sample was then split into two fractions: about 2 g for elemental analysis by ICP-OES and ICP-MS, and about 20 g for analysis of ^{36}Cl by accelerator mass spectrometry (AMS). Chlorine was extracted and purified from the 125–250 μm fraction of leached samples and precipitated as AgCl using modified version of procedures developed by Stone *et al.* (1996). Samples were processed together with a full chemistry blanks.

$^{10}\text{Be}/^9\text{Be}$, $^{36}\text{Cl}/^{35}\text{Cl}$ and $^{36}\text{Cl}/^{37}\text{Cl}$ ratios were measured using the 5MW pelletron at SUERC (Xu *et al.*, 2010, Wilcken *et al.* 2013) and normalised to NIST SRM4325 with a $^{10}\text{Be}/^9\text{Be}$ ratio of 2.79×10^{-11} (Nishiizumi *et al.*, 2007), and Z93-0005 (PRIME Lab, Purdue) with a $^{36}\text{Cl}/\text{Cl}$ ratio of 1.2×10^{-12} . Cosmogenic nuclide concentrations include a blank correction of 3-14% for ^{10}Be and 1-5% for ^{36}Cl (Table 3). The uncertainties in the cosmogenic nuclide concentrations include the AMS

counting statistics and scatter uncertainties from sample, procedural blank, and standards measurements.

Age calculation and filtering

The cosmogenic ^{10}Be ages were calculated using two methods. First, ages were determined using version 2.3 of the online calculators formerly known as CRONUS-Earth ^{10}Be - ^{26}Al exposure age calculators (Balco *et al.*, 2008: http://hess.ess.washington.edu/math/al_bev_23/al_be_multiple_v23.php) using the independently-constrained Loch Lomond production rate (LLPR; nominal production rate 4.00 ± 0.18 atoms $\text{g}^{-1} \text{a}^{-1}$; Fabel *et al.*, 2012), the default production rate employed in the BRITICE-CHRONO dating programme. Exposure ages were based on the time-dependent Lm scaling (Lal, 1991; Stone, 2000) and assume 1 mm ka^{-1} of post-depositional surface erosion (cf. André, 2002; Nicholson, 2009; Larsen *et al.*, 2012).

We also calculated cosmogenic ^{10}Be exposure ages using the CRONUScalc program version 2.0 (Marrero *et al.*, 2016a) and the default global production rate of 3.92 atoms $\text{g}^{-1} \text{a}^{-1}$ for Lm scaling (Borchers *et al.*, 2016), again assuming an erosion rate of 1 mm ka^{-1} . Both production rates agree within $\pm 1\sigma$ uncertainties with the range of production rates determined for other high latitude sites in the northern hemisphere (Phillips *et al.*, 2016).

^{36}Cl ages are presented as determined using the CRONUScalc calculator which uses production rates of 56 ± 4.1 at $^{36}\text{Cl} (\text{g Ca})^{-1} \text{a}^{-1}$ for Ca spallation, 155 ± 11 at $^{36}\text{Cl} (\text{g K})^{-1} \text{yr}^{-1}$ for K spallation and 759 ± 180 neutrons $(\text{g air})^{-1} \text{a}^{-1}$ (Marrero *et al.*, 2016b). Table 3 presents the ^{10}Be and ^{36}Cl data and exposure ages with associated uncertainties for both methods of calculation.

Within-site consistency of ages was tested using the reduced Chi-square statistic (χ^2_{R}) (Bevington and Robinson, 2003). Where the χ^2_{R} value for a sample of ages from a single site

1
2
3 288 exceeds the critical value at the 95% level, it was inferred that geological uncertainty contributed
4
5 289 to the observed age scatter. In such cases outlier ages were manually removed until a χ^2_R value less
6
7 290 than the critical value was obtained; the remaining ages were regarded as consistent with and
8
9
10 291 representative of a single age population, with age scatter being due to measurement error alone
11
12 292 (Balco, 2011; Applegate *et al.*, 2012; Small and Fabel, 2016; Small *et al.*, 2017a). For sites having
13
14 293 two or more internally consistent ages the uncertainty-weighted mean was determined and is
15
16 294 regarded as providing the best estimate exposure age for the site. As with the legacy TCN ages
17
18 295 discussed above, we cite the ^{10}Be weighted mean ages determined with the LLPR first, followed
19
20 296 by the equivalent ages calculated with CRONUScale in brackets.
21
22
23
24 297

25
26 298 **Results**
27

28
29 299 The 20 new TCN surface exposure ages and uncertainty-weighted mean values for internally
30
31 300 consistent ages for each site are given in Table 3 and Fig. 3. These ages are assessed below in
32
33 301 relation to published ages for the region (Table 1).
34

35
36 302 The ages for Rosguill and Malin Head, on the north coast of Donegal, complement
37
38 303 published deglacial age estimates for the northern sites of Aran Island, Bloody Foreland and
39
40 304 Corvish. A mean value was not calculated for Malin Head because the three samples failed to yield
41
42 305 an acceptable χ^2_R value, due to their wide age scatter ($\sim 25.5\text{--}20.7$ ka and $25.2\text{--}20.2$ ka). However,
43
44 306 sample MH-03 yielded an age of 20.7 ± 1.2 ka (20.2 ± 1.7 ka), reasonably consistent with the TCN
45
46 307 mean ages of 21.7 ± 1.1 ka (21.0 ± 1.7 ka) for Aran Island, 21.8 ± 1.2 ka (21.1 ± 1.8 ka) for Bloody
47
48 308 Foreland, and the mean age of 18.8 ± 0.9 ka (18.6 ± 1.5 ka) for Rosguill; in addition, the MH-03 age
49
50 309 is consistent with the minimum deglaciation ^{14}C age of 20.68 ± 0.16 cal ka BP for Corvish, and it
51
52 310 therefore provides the best fit age of the three Malin Head ages. Furthermore, Malin Head is the
53
54
55
56
57
58
59
60

311 most easterly of our sites and is unlikely to have been deglaciated before the more westerly sites.

312 The other two Malin Head samples are considered to be compromised by nuclide inheritance.

313 The mean age of 16.7 ± 0.9 ka (16.2 ± 1.4 ka) for the Poisoned Glen boulder limit in north
314 Donegal is statistically indistinguishable from the mean age of 17.8 ± 0.9 ka (17.6 ± 1.4 ka) obtained
315 from ice-plucked bedrock on Errigal, 2.2 km north and 350 m higher. Together these two sites
316 indicate that the northern mountains were largely deglaciated by ~ 18 -17 ka.

317 In south Donegal, the consistent exposure ages obtained from three boulders from
318 Glencolumbkille yield a mean age of 16.7 ± 0.8 ka (16.4 ± 1.4 ka). Two legacy samples from this
319 location had returned ages of 17.9 ± 0.9 ka (17.4 ± 1.4 ka) and 19.8 ± 1.0 ka (19.1 ± 1.5 ka). The former
320 age is statistically indistinguishable from the three new ages, and collectively all four ages produce
321 an uncertainty weighted mean age of 17.2 ± 0.8 ka (16.8 ± 1.3 ka) ($\chi^2_R = 1.75$ and 1.64 respectively).
322 The latter age possibly reflects the influence of nuclide inheritance. The Glencolumbkille ages are
323 also consistent with three legacy samples from rockslide runout debris on Slieve League (7 km SE
324 of Glencolumbkille) that yielded a mean minimum age for deglaciation of 17.1 ± 0.9 ka (16.9 ± 1.4
325 ka).

326 Four samples from dolerite boulders at Kilcar gave ^{36}Cl exposure ages ranging from
327 18.1 ± 1.7 ka to 42.0 ± 6.0 ka. Samples KC-02, -03 and -04 returned ages that pre-date the LGM and
328 are probably compromised by nuclide inheritance. The other sample (KC-01: 18.1 ± 1.7 ka) is
329 consistent with the wider geochronological evidence for the timing of deglaciation in south
330 Donegal and north Mayo.

331 The four samples obtained from boulders on low ground (~ 150 m) at the foot of the Blue
332 Stack Mountains yielded consistent ($\chi^2_R < 1.0$) ages ranging from 15.6 ± 0.9 ka (15.6 ± 1.4 ka) to
333 14.4 ± 0.8 ka (14.5 ± 1.3 ka), and an uncertainty-weighted mean age of 14.8 ± 0.7 ka (14.8 ± 1.2 ka).
334 These results imply deglaciation of all low ground at the head of Donegal Bay before ~ 14.7 ka.

1
2
3 335 They also suggest that ice persisted much later in the Blue Stack Mountains than in the Derryveagh
4
5 336 Mountains of northern Donegal, where the available dating evidence suggests deglaciation of the
6
7 337 Errigal col at ~17.8 ka and ice withdrawal from the Poisoned Glen boulder limit at ~16.7 ka.
8
9
10 338

11
12 339 **Discussion**
13
14

15 340 In conjunction with the published legacy ages discussed earlier, the new ages presented here
16
17 341 provide spatially-consistent constraints on the timing of deglaciation in Donegal (Fig. 3). Below
18
19 342 we discuss deglaciation of northern and southern Donegal separately, because during the ILGM
20
21 343 northern ice fed the Hebrides/Malin Sea Ice Stream that drained ice from western Scotland across
22
23 344 the Malin Shelf, whereas southern ice contributed to the Donegal Bay Ice Stream that drained ice
24
25 345 from the Irish Midlands to the shelf edge through Donegal Bay.
26
27
28

29 346
30
31 347 *Deglacial chronology of north Donegal*
32

33 348 The TCN ages relating to deglaciation of Aran Island (mean = 21.7±1.1 ka (21.0±1.7 ka)), Bloody
34
35 349 Foreland (mean = 21.8±1.2 ka (21.1±1.8 ka)), and Malin Head (a single TCN age of 20.7±1.2 ka
36
37 350 (20.2±1.7 ka)), together with the oldest ¹⁴C age from Corvish (20.68±0.16 cal ka BP) indicate
38
39 351 progressive eastward retreat of the ice margin along the northern coast of Donegal between
40
41 352 ~21.8 ka and ~20.7 ka. The Bloody Foreland and Aran Island ages imply that decoupling of ice
42
43 353 sourced in Donegal from the Scottish-sourced Hebridean Ice Stream commenced within the
44
45 354 interval ~22–21 ka; this is slightly earlier than previous estimates, which have placed initial
46
47 355 disengagement of these two ice masses after ~21 ka (Small *et al.*, 2017). The single Malin Head
48
49 356 TCN age and the oldest ¹⁴C age at Corvish indicate that separation of Scottish-sourced ice and
50
51 357 Donegal-sourced ice was complete by ~20.7 ka, implying that by this time a marine embayment
52
53
54
55
56
57
58
59
60

extended eastward along the north coast of Donegal, separating ice flowing north and northeast from the Donegal Ice Centre from the retreating Hebrides/Malin Sea Ice Stream.

The timing of ice retreat inland towards the Derryveagh Mountains of northern Donegal is provided by the TCN ages obtained for the two sites in the heart of this range, at ~420 m OD on Errigal col (mean = 17.8 ± 0.9 ka (17.6 ± 1.4 ka)) and, 2.2 km to the south, a low-level site (~74 m OD) at the mouth of the Poisoned Glen (mean = 16.7 ± 0.9 ka (16.2 ± 1.4 ka)). Although these two ages are statistically indistinguishable within uncertainties (Fig. 4), the difference between them may imply exposure of Errigal col by downwasting ice several centuries prior to retreat of ice in the Poisoned Glen. Irrespective of whether or not this was the case, the deglaciation ages for both sites imply that ~3000–5000 years elapsed between deglaciation of Aran Island and Bloody Foreland and deglaciation of the Derryveagh Mountains (Fig. 3). A further implication is that net ice margin retreat rates were extremely slow. The Errigal col and Poisoned Glen sites lie respectively 15 km and 17 km SE of the Bloody Foreland site; if the mean deglaciation ages for these sites are representative, then the net ice-margin retreat rate from Bloody Foreland to both sites was ~ 4 m a^{-1} ; taking the associated uncertainties into account suggests that net retreat rate is unlikely to have exceeded 5 m a^{-1} , and may have been as low as 3 m a^{-1} . By contrast, assuming that the ice margin began to retreat from the shelf edge within the interval 26.3 ka to 24.8 ka (Ó Cofaigh *et al.*, 2018), the implied net rate of offshore ice-margin retreat from the shelf break to Bloody Foreland falls within the range ~ 19.2 – 33.3 m a^{-1} . Ó Cofaigh *et al.* (2018) inferred that ice-sheet retreat from the shelf edge was initiated by calving associated with high sea levels induced by glacio-isostatic depression rather than changing climate, and the marked slowing of retreat after the ice margin had become land-based in northern Donegal appears consistent with this interpretation: the inferred slow net retreat rates of Donegal ice in this sector over the period ~21

381 ka to ~17 ka suggest that the retreating ice was close to equilibrium with prevailing climate, and
382 experienced only a slight net negative mass balance during this period.

383 Averaged net retreat rates, however, may obscure oscillations of the ice margin, with
384 periods of retreat alternating with limited readvances. At present there is dated stratigraphic
385 evidence for only one such readvance, at Corvish, near the head of Trawbreaga Bay (Fig. 3).
386 Readvance occurred over a distance of at least 5 km according to McCabe and Clark (2003). At
387 this site, the youngest radiocarbon age obtained for *Elphidium clavatum* tests in deformed marine
388 silts (18.32 ± 0.18 cal ka BP) and a single age for *E. clavatum* tests in overlying undeformed silts
389 (17.06 ± 0.18 cal ka BP) have been interpreted by McCabe and Clark (2003) as bracketing the
390 timing of readvance of the ice margin on the northern coast of Donegal. They placed the timing of
391 this readvance at ~18 ka, though the dating evidence appears consistent with readvance of the ice
392 margin at any time within the interval ~18.5–16.9 ka. On the assumption that the Corvish readvance
393 occurred at ~18 ka, McCabe *et al.* (2007) suggested that it correlates with the Clogher Head
394 Readvance (CHR) in NE Ireland, though reinterpretation of the stratigraphic and dating evidence
395 indicates that the CHR was a short-lived event that peaked rather earlier, at ~18.4 ka (Ballantyne
396 and Ó Cofaigh, 2017). Thus although the two readvances may be coeval and represent a regional-
397 scale event that occurred in response to climatic forcing (McCabe *et al.*, 2007; J. Clark *et al.*, 2012)
398 it is equally feasible that they occurred at different times and represent localised oscillations of the
399 ice margin (C.D. Clark *et al.*, 2012). The TCN mean age of 18.8 ± 0.9 ka (18.6 ± 1.5) indicative of
400 the timing of deglaciation at Rosguill predates the bracketing ages for the readvance at Corvish
401 (~18.5–16.9 ka), but because of the uncertainties associated with the TCN age we cannot not
402 preclude the possibility that the Rosguill site was reoccupied by glacier ice during the same
403 readvance event.

The TCN mean age for the Poisoned Glen (16.7 ± 0.9 ka (16.2 ± 1.4 ka) implies that ice persisted in the Derryveagh Mountains until ~ 17 – 16 ka, but the ^{14}C age of 15.38 ± 0.12 cal. ka BP from Lough Nadourcan (Watson *et al.*, 2010) suggests that ice had disappeared from low ground surrounding these mountains before the rapid warming associated with the onset of the Lateglacial Interstadial at ~ 14.7 ka. Valley heads, cirques and plateaus in the Derryveagh Mountains may, however, have retained ice until early in the interstadial, as appears to have been the case for the Blue Stack Mountains of south Donegal (see below).

Deglacial chronology of south Donegal and Donegal Bay

As noted earlier, the Donegal Bay Moraine (DBM) that crosses outer Donegal Bay (Fig. 3) represents the limit of a readvance of ice fed from Donegal Bay. Radiocarbon ages of 20.24 ± 0.24 cal ka BP and 17.92 ± 0.16 cal ka BP obtained for foraminifera retrieved respectively from the distal and proximal sides of the moraine constrain its age (Ó Cofaigh *et al.*, 2018). This broad interval encompasses the timing of both the readvance at Corvish in northern Donegal and that of the Clogher Head Readvance in NE Ireland (McCabe and Clark, 2003; McCabe *et al.*, 2007; J. Clark *et al.*, 2012; Ballantyne and O Cofaigh, 2017), but the resolution of the dating evidence is inadequate to establish contemporaneity. The position and alignment of the DBM suggests that the sites at Belderg Pier and Fiddauntawnoneen on the south coast of outer Donegal Bay lay outside the readvance, and the radiocarbon ages of ~ 20 – 19 cal ka BP obtained by McCabe *et al.* (1986, 2005) for *in situ* marine fauna within glaci-marine sediments at these sites (Table 1) are consistent with this interpretation (Fig. 3). Conversely the aggregated TCN mean age for Glencolumbkille in SE Donegal (17.2 ± 0.8 ka (16.8 ± 1.3 ka)), the single TCN age for Kilcar (18.1 ± 1.7 ka), and the minimum deglaciation age represented by postglacial rockslide debris at nearby Slieve League (17.1 ± 0.9 ka (16.9 ± 1.4 ka)) suggest that southern Donegal lay within the limits of the readvance

1
2
3 428 that produced the DBM. Similarly, the five ‘younger’ TCN ages
4
5 429 (mean = $16.6 \pm 0.1.0$ ka (16.7 ± 1.5 ka)) reported by Clark *et al.* (2009) for the Twanywaddyduff
6
7 430 moraine system of the northern Ox Mountains (Fig. 3; Table 1) indicate persistence of ice cover
8
9 431 along the inner part of Donegal Bay after ~17 ka. Collectively, these two sets of ages suggest that
10
11 432 much or all of Donegal Bay continued to support ice cover as late as ~17 ka, though it is possible
12
13 433 that a calving margin along the axis of the bay led to development of an ice-free marine corridor
14
15 434 between its northern and southern shores.
16
17
18

19 435 The TCN mean age of 14.8 ± 0.7 ka (14.8 ± 1.2 ka) from the southern flanks of the Blue Stack
20
21 436 Mountains suggests that Donegal Bay had become ice free by ~15 ka but that ice still occupied
22
23 437 mountain valleys and cirques near the head of the bay (Fig. 3). The implication of this age is that
24
25 438 mountain ice probably persisted for some time following the onset of the Lateglacial Interstadial
26
27 439 at ~14.7 ka. The Blue Stack ages are the youngest ages for deglaciation hitherto reported for Ireland
28
29 440 (*cf.* Ballantyne and Ó Cofaigh, 2017), suggesting that these and possibly other mountains in NW
30
31 441 Ireland supported the last remnants of the last Irish Ice Sheet prior to complete disappearance of
32
33 442 glacier ice under the warmer conditions of the Lateglacial Interstadial.
34
35
36

37 443 The sampling sites at Glencolumbkille and the Blue Stack Mountains are separated by a
38
39 444 distance of 41 km. The mean TCN ages for these two sites imply that net ice margin retreat between
40
41 445 these two sites occurred over ~2,200 years, implying a net retreat rate of ~19 m yr⁻¹; taking the
42
43 446 associated age uncertainties into account implies that net retreat rate of the ice margin along the
44
45 447 northern shore of Donegal Bay fell within the range 12–24 m a⁻¹, markedly faster than the net rate
46
47 448 inferred above (3–5 m a⁻¹) for ice retreat inland from Bloody Foreland to the Derryveagh
48
49 449 Mountains. Although subject to the same caveat (that retreat may have been interrupted by one or
50
51 450 more ice margin readvances), there is neither morphological nor seismostratigraphic evidence for
52
53 451 later readvances of the ice margin as it retreated eastward from the DBM: Ó Cofaigh *et al.* (2018)
54
55
56
57
58
59
60

noted that the sediment cover east of the moraine comprises undeformed, acoustically-stratified conformable glacimarine sediments overlain by postglacial marine deposits. For comparison, the average net rate calculated by Ó Cofaigh *et al.* (2018) for ice-margin retreat from the shelf edge to the DBM is 11.2–14.0 m a⁻¹, though they considered that this probably incorporated moderately rapid retreat at a minimum rate of 35.7 m a⁻¹ from the shelf edge to mid-shelf, followed by much slower oscillatory retreat at a net rate of 5.5 m a⁻¹ between the mid-shelf and the DBM.

Wider implications

Collectively, the chronological data reported above indicate a marked contrast in both the timing and rate of net ice margin recession of land-based ice in northern Donegal (~21–17 ka) and retreat of the ice margin in southern Donegal adjacent to Donegal Bay (~17–15 ka). This contrast suggests that the timing of ice-margin retreat was at least partly conditioned by the relationship between ice fed from the Donegal Ice Centre and adjacent ice streams. Northern Donegal lay in an inter-ice-stream location, between the Hebrides/Malin Sea Ice Stream to the north and the Donegal Bay Ice Stream to the south, and here the early (~22–21 ka) decoupling of Donegal ice from the extended Hebrides/Malin Sea Ice Stream appears to have created an ice-free marine embayment along the north coast of Donegal, so that ice flowing from the Donegal Ice Centre was effectively unconstrained, and subsequently retreated gradually in response to a slight net negative mass balance. In contrast, the south coast of Donegal lay near the axis of the Donegal Bay Ice Stream, which was fed not only by ice from the Donegal Ice Centre, but also by ice from the Irish Midlands. Following initial rapid retreat, the oscillating margin of the Donegal Bay Ice Stream retreated slowly from mid-shelf to the DBM (Ó Cofaigh *et al.*, 2018), so that ice cover persisted over south Donegal until ~17 ka, after which it retreated to the footslopes of the Blue Stack Mountains. This contrast in behaviour implies that different dynamics apply to extended marine-based ice streams,

1
2
3 476 which are sensitive to changes in sea level, confinement and bed slope (Smedley *et al.*, 2017; Ó
4
5 477 Cofaigh *et al.*, 2018; Small *et al.*, 2018) and land-based ice masses in inter-ice-stream locations,
6
7
8 478 which respond mainly to changes in climate inputs.
9

10 479 Evidence for marked slowing of ice-margin retreat as the shrinking BIIS stabilised at or
11
12 480 near the present coastline is not limited to northern Donegal. TCN ages reported by Small *et al.*
13
14 481 (2017) for the Sea of the Hebrides to the west of Scotland suggest that termination of ice streaming
15
16 482 after ~20.6 ka was succeeded by a ~3000–4000 year interval during which the ice margin
17
18 483 experienced oscillatory net retreat of only 50–70 km as it became progressively land-based
19
20 484 amongst the islands of the Inner Hebrides. The slowing of ice margin retreat in this area coincides
21
22 485 closely with the period of very gradual ice-margin recession in northern Donegal.
23
24
25

26 486 The mean exposure age of the samples from low ground (~150 m) at the foot the Blue Stack
27
28 487 Mountains (14.8 ± 0.7 ka (14.8 ± 1.2 ka)) represents the youngest age for the timing of ice-sheet
29
30 488 deglaciation hitherto reported for Ireland and implies that by ~14.7 ka the Donegal Ice Centre had
31
32 489 shrunk to a small ice cap or transection complex centred on high ground. An analogous situation
33
34 490 occurred in SW Scotland, 240 km to the east, where seven (recalibrated) TCN ages indicate that
35
36 491 only fragmented upland remnants of the Galloway Hills Ice Centre remained by ~15.1 ka
37
38 492 (Ballantyne *et al.*, 2013a). The Galloway Hills TCN ages are statistically indistinguishable from
39
40 493 the Blue Stack ages, and both confirm that almost all ice retreat occurred under stadial conditions
41
42 494 prior to ~14.7 ka. In both areas it is unlikely that remnant glacier ice survived subsequent rapid
43
44 495 warming, when mean July temperatures inferred from subfossil chironomid assemblages rose
45
46 496 rapidly by 5–6°C (Brooks and Birks, 2000; Lang *et al.*, 2010; Watson *et al.*, 2010, Van Asch *et al.*,
47
48 497 2012). A more general implication is that all of Ireland and southern Scotland were probably
49
50
51
52
53 498 completely deglaciated early in the Lateglacial Interstade. For the British Isles as a whole, present
54
55
56
57
58
59
60

evidence suggests that remnants of the BIIS survived the interstade (if at all) only in the Highlands of Scotland (Finlayson *et al.* 2011; Ballantyne and Small, 2018).

Conclusions

1. Twenty new TCN ages obtained for sites in northern and southern Donegal complement (and are broadly consistent with) previously-published TCN and radiocarbon ages, and reveal marked contrasts in the timing and rate of deglaciation in northern and southern Donegal.
2. The TCN ages for northern Donegal indicate decoupling of ice fed from the Donegal Ice Centre from the Hebrides/Malin Shelf Ice Stream and associated development of a marine embayment north of Donegal by ~22–21 ka. Conversely, the new TCN ages for south Donegal confirm that ice persisted in much or all Donegal Bay and covered southwest Donegal as late as ~17 ka.
3. In northern Donegal our TCN data imply very gradual ice-margin retreat inland towards mountain source areas at a net rate of $4 \pm 1 \text{ m a}^{-1}$; by comparison, the inferred net rate of ice margin retreat from SW Donegal to the foothills of the Blue Stack Mountains near the head of Donegal Bay averaged $18 \pm 6 \text{ m a}^{-1}$.
4. We suggest that the above contrast in timing and rate of ice retreat reflect differences in location relative to those of major ice streams. Northern Donegal occupied an inter-ice-stream location, and after early decoupling of Donegal-sourced ice from the Hebrides/Malin Sea Ice Stream the former was unconstrained and retreated mainly in response to changes in climatic inputs. Conversely, southern Donegal lay near to the axis of the Donegal Bay Ice Stream, which occupied (or reoccupied) much of Donegal Bay as late as ~18 ka, delaying deglaciation along the southern coast of Donegal until after ~17 ka.
5. A mean TCN age of $\sim 14.8 \pm 0.7 \text{ ka}$ ($14.8 \pm 1.2 \text{ ka}$) obtained for the footslopes of the Blue Stack mountains in southern Donegal is the youngest deglacial age hitherto reported for Ireland, and

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

implies that shortly before the onset of rapid warming at the beginning of the Lateglacial Interstade (~14.7 ka) the Donegal Ice Centre had shrunk to a small icecap or icefield of very limited extent, and probably disappeared completely during the early part of the interstade. This date also confirms that virtually all of the retreat of the Irish Ice Sheet occurred under stadial conditions prior to the onset of interstadial warming.

Acknowledgements

This work was supported by the UK Natural Environment Research Council consortium grant: BRITICE-CHRONO NE/J009768/1. The TCN analyses were undertaken at the NERC Cosmogenic Isotope Analysis Facility (allocation 9155/1014) Staff at the Scottish Universities Environmental Research Centre AMS Laboratory, East Kilbride, are thanked for ¹⁰Be and ³⁶Cl isotope measurements.

References

- André M-F. 2002. Rates of postglacial rock weathering on glacially scoured outcrops (Abisko-Riksgränsen area, 68 °N). *Geografiska Annaler* **84A**: 139-150.
- Applegate PJ, Urban NM, Keller K *et al.* 2012. Improved moraine age interpretations through explicit matching of geomorphic process models to cosmogenic nuclide measurements from single landforms. *Quaternary Research* **77**: 293-304.
- Benetti S, Dunlop P, Ó Cofaigh C. 2010. Glacial and glacially-related features on the continental margin of northwest Ireland mapped from marine geophysical data. *Journal of Maps* **2010**: 14-29.
- Balco G. 2011. Contributions and unrealized potential contributions of cosmogenic-nuclide exposure dating to glacier chronology, 1990-2010. *Quaternary Science Reviews* **30**: 3-27.
- Balco G, Stone JO, Lifton NA, Dunai TJ. 2008. A complete and easily accessible means of calculating surface exposure ages or erosion rates from ^{10}Be and ^{26}Al measurements. *Quaternary Geochronology* **3**: 174-195.
- Ballantyne CK, Ó Cofaigh C. 2017. The last Irish Ice Sheet: extent and chronology. In: Coxon P, Mitchell F, McCarron S. (eds), *Advances in Irish Quaternary Studies*. Atlantic Press, Paris, 101-149.
- Ballantyne CK, Small D. 2018. The Last Scottish Ice Sheet. *Earth and Environmental Science Transactions of the Royal Society of Edinburgh*. DOI: 10.1017/S1755691018000038
- Ballantyne CK, McCarroll D, Stone JO. 2007. The Donegal ice dome, northwest Ireland: dimensions and chronology. *Journal of Quaternary Science* **22**: 773-783.
- Ballantyne CK, McCarroll D, Stone JO. 2011. Periglacial trimlines and the extent of the Kerry-Cork Ice Cap, SW Ireland. *Quaternary Science Reviews* **30**: 3834-3845.
- Ballantyne CK, Rinterknecht V, Gheorghiu DM. 2013a. Deglaciation chronology of the Galloway Hills Ice Centre, southwest Scotland. *Journal of Quaternary Science* **28**: 412-420.
- Ballantyne CK, Wilson P, Schnabel C, Xu S. 2013b. Lateglacial rock slope failures in north-west Ireland: age causes and implications. *Journal of Quaternary Science* **28**: 789-802.
- Bevington PR, Robinson DK. 2003 *Data reduction and error analysis for the physical sciences*. McGraw-Hill, New York.
- Borchers B, Marrero S, Balco G *et al.* 2016. Geological calibration of spallation production rates in the CRONUS-Earth project. *Quaternary Geochronology* **31**: 188-198.
- Bowen DQ, Rose J, McCabe AM, Sutherland DG. 1986. Correlation of Quaternary glaciations in England, Ireland, Scotland and Wales. *Quaternary Science Reviews* **5**: 299-340.

- Bronk Ramsey C. 2009. Bayesian analysis of radiocarbon dates. *Radiocarbon* **51**: 337-360.
- Brooks SJ, Birks HJB. 2000. Chironomid-inferred Late-glacial temperatures at Whitrig Bog, southeast Scotland. *Journal of Quaternary Science* **15**: 759-764.
- Charlesworth JK. 1924. The glacial geology of north-west Ireland. *Proceedings of the Royal Irish Academy* **36B**: 174-314.
- Child D, Elliott G, Mifsud C *et al.* 2000. Sample processing for earth science studies at ANTARES. *Nuclear Instruments and Methods in Physics Research B* **172**: 856-860.
- Clark J, McCabe AM, Schnabel C *et al.* 2009a. ^{10}Be chronology of the last deglaciation of County Donegal, northwestern Ireland. *Boreas* **38**: 111-118.
- Clark J, McCabe AM, Schnabel C *et al.* 2009b. Cosmogenic ^{10}Be chronology of the last deglaciation of western Ireland, and implications for sensitivity of the Irish Ice Sheet to climate change. *Geological Society of America Bulletin* **121**: 3-16.
- Clark CD, Hughes ALC, Greenwood, SL *et al.* 2012. Pattern and timing of retreat of the last British-Irish Ice Sheet. *Quaternary Science Reviews* **44**: 112-146.
- Clark J, McCabe AM, Bowen DQ, Clark PU. 2012. Response of the Irish Ice Sheet to abrupt climate change during the last deglaciation. *Quaternary Science Reviews* **35**: 100-115.
- Clark PU, Dyke, AS, Shakun JD *et al.* 2009. The Last Glacial Maximum. *Science* **325**: 710-714.
- Colhoun EA. 1973. Two Pleistocene sections in south-eastern Donegal and their relation to the last glaciation of the Glengesh plateau. *Irish Geography* **6**: 594-609.
- Cullen C. 2013. *Deciphering the geomorphic and sedimentary record of the last Irish Ice Sheet in NW Donegal, Ireland: implications for glacial dynamics and decay configurations*. Unpublished PhD thesis, National University of Ireland, Galway.
- Dunlop P, Shannon R, McCabe M *et al.* 2010. Marine geophysical evidence for ice sheet extension and recession on the Malin Shelf: new evidence for the western limits of the British Irish Ice Sheet. *Marine Geology* **276**: 86-90.
- Dury GH. 1957. A glacially breached watershed in Donegal. *Irish Geography* **3**: 171-180.
- Dury GH. 1958. Glacial morphology of the Blue Stack area, Donegal. *Irish Geography* **3**: 242-253.
- Dury GH. 1964. Aspects of the geomorphology of Slieve League Peninsula, Donegal. *Proceedings of the Geologists' Association* **75**: 445-459.
- Fabel D, Ballantyne CK, Xu S. 2012. Trimlines, blockfields, mountain-top erratics and the vertical dimensions of the last British-Irish ice sheet in NW Scotland. *Quaternary Science Reviews* **55**: 91-102.

- Finlayson A, Golledge N, Bradwell T, Fabel D. 2011. Evolution of a Lateglacial mountain icecap in northern Scotland. *Boreas* **40**: 536–554.
- Greenwood SL, Clark CD. 2009a. Reconstructing the last Irish Ice Sheet 1: changing flow geometries and ice flow dynamics deciphered from the glacial landform record. *Quaternary Science Reviews* **28**: 3085–3100.
- Greenwood SL, Clark CD. 2009b. Reconstructing the last Irish Ice Sheet 2: a geomorphologically-driven model of ice-sheet growth, retreat and dynamics. *Quaternary Science Reviews* **28**: 3101–3123.
- Hubbard A., Bradwell T, Golledge N *et al.* 2009. Dynamic cycles, ice streams and their impact on the extent, chronology and deglaciation of the British–Irish ice sheet. *Quaternary Science Reviews* **28**: 758–776.
- Knight J. 2003. Geomorphic and sedimentary evidence for patterns of late Midlandian ice retreat in the Tempo valley, north-central Ireland. *Irish Geography* **36**: 127–144.
- Knight J. 2012. The last glaciation of Aran Island and Cruit Island, Co. Donegal, north-west Ireland. *Irish Journal of Earth Sciences* **30**: 49–58.
- Knight J, McCabe AM. 1997. Drumlin evolution and ice sheet oscillations along the NE Atlantic margin, Donegal bay, western Ireland. *Sedimentary Geology* **111**: 57–72.
- Kohl CP, Nishiizumi K. 1992. Chemical isolation of quartz for measurement of in-situ -produced cosmogenic nuclides. *Geochimica et Cosmochimica Acta* **56**: 3583–3587.
- Lal D. 1991. Cosmic ray labelling of erosion surfaces: in situ nuclide production rates and erosion models. *Earth and Planetary Science Letters* **104**: 424–439.
- Lang B, Brooks SJ, Bedford A *et al.* 2010. Regional consistency in Lateglacial chironomid-inferred temperatures from five sites in north-west England. *Quaternary Science Reviews* **29**: 1528–1538.
- Larsen NK, Linge H, Håkansson L, Fabel D. 2012. Investigating the last deglaciation of the Scandinavian Ice Sheet in southwest Sweden with ^{10}Be exposure dating. *Journal of Quaternary Science* **27**: 211–220.
- Long CB, McConnell BJ. 1997. *Geology of North Donegal*. Geological Survey of Ireland, Dublin.
- Long CB, McConnell BJ. 1999. *Geology of South Donegal*. Geological Survey of Ireland, Dublin.
- Marrero SM, Phillips FM, Borchers B *et al.* 2016a. Cosmogenic nuclide systematics and the CRONUScale program. *Quaternary Geochronology* **31**: 160–187.
- Marrero SM, Phillips FM, Caffee MW, Gosse JC. 2016b. CRONUS-Earth cosmogenic ^{36}Cl calibration. *Quaternary Geochronology*, **31**: 199–219.

- McCabe AM. 1985. Glacial geomorphology. In: Edwards KJ, Warren WP. (eds), *The Quaternary history of Ireland*. Academic Press, London, 67-93.
- McCabe AM. 1995. Quaternary geology. In: Wilson P. (ed.), *North-west Donegal*. Field Guide No. 19, Irish Association for Quaternary Studies, Dublin, 15-20.
- McCabe AM, Clark PU. 2003. Deglacial chronology from County Donegal, Ireland: implications for deglaciation of the British-Irish ice sheet. *Journal of the Geological Society, London*, **160**: 847-855.
- McCabe AM, Bowen DQ, Penney DN. 1993. Glaciomarine facies from the western sector of the last British Ice Sheet, Malin Beg, County Donegal, Ireland. *Quaternary Science Reviews* **12**: 35-45.
- McCabe AM, Clark PU, Clark J. 2005. AMS ^{14}C dating of deglacial events in the Irish Sea Basin and other sectors of the British-Irish ice sheet. *Quaternary Science Reviews* **24**: 1673-1690.
- McCabe AM, Clark PU, Clark J, Dunlop P. 2007. Radiocarbon constraints on readvances of the British-Irish Ice Sheet in the northern Irish Sea Basin during the last deglaciation. *Quaternary Science Reviews* **26**: 1204-1211.
- McCabe AM, Haynes JR, McMillan NF. 1986. Late Pleistocene tidewater glaciers and glaciomarine sequences from north County Mayo, Republic of Ireland. *Journal of Quaternary Science* **1**: 73-84.
- Nicholson DT. 2009. Holocene microweathering rates and processes on ice-eroded bedrock, Rødal area, Hardangervidda, southern Norway. In: Knight J, Harrison S. (eds), *Periglacial and Paraglacial Processes and Environments*. Geological Society, London, Special Publications **320**: 29-49.
- Nishiizumi K, Imamura M, Caffee MW *et al.* 2007. Absolute calibration of ^{10}Be AMS standards. *Nuclear Instruments and Methods in Physics Research B* **258**: 403-413.
- Ó Cofaigh C, Dunlop P, Benetti S. 2012. Marine geophysical evidence for Late Pleistocene ice sheet extent and recession off northwest Ireland. *Quaternary Science Reviews* **44**: 147-159.
- Ó Cofaigh C, Weilbach K, Lloyd JM *et al.* 2018. Early deglaciation of the British-Irish Ice Sheet on the Atlantic shelf northwest of Ireland driven by glacioisostatic depression and high relative sea level. *Quaternary Science Reviews* in revision.
- Phillips FM, Argento DC, Balco G *et al.* 2016. The CRONUS earth project: a synthesis. *Quaternary Geochronology* **31**: 119-154.
- Rasmussen SO, Bigler M, Blockley SP *et al.* 2014. A stratigraphic framework for abrupt climatic changes during the Last Glacial period based on three synchronized Greenland ice-core records: refining and extending the INTIMATE event stratigraphy. *Quaternary Science Reviews* **106**: 14-28.

- Reimer P, Bard E, Bayliss A *et al.* 2013. Intercal13 and Marine13 radiocarbon age calibration curves 0-50,000 years cal BP. *Radiocarbon* **55**: 1869-1887.
- Sejrup HP, Hjelstuen BO, Dahlgren KIT *et al.* 2005. Pleistocene glacial history of the NW European continental margin. *Marine and Petroleum Geology* **22**: 1111-1129.
- Sellier D. 1995. Derryveagh Mountains – glacial evidence. In: Wilson P. (ed.), *North-west Donegal*. Field Guide No. 19, Irish Association for Quaternary Studies, Dublin, 57-58.
- Small D, Fabel D. 2016. Response to Bromley *et al.* “Comment on ‘Was Scotland deglaciated during the Younger Dryas?’ by Small and Fabel (2016). *Quaternary Science Reviews* **152**: 206-208.
- Small D, Clark CD, Chiverrell RC *et al.* 2017a. Devising quality assurance procedures for assessment of legacy geochronological data relating to deglaciation of the last British-Irish Ice Sheet. *Earth-Science Reviews* **164**: 232-250.
- Small D, Benetti S, Dove D *et al.* 2017b. Cosmogenic exposure age constraints on deglaciation and flow behaviour of a marine-based ice stream in western Scotland, 21-16 ka. *Quaternary Science Reviews* **167**: 30-46.
- Small D, Smedley RK, Chiverrell RC *et al.* Trough geometry was a greater influence than *climate-ocean forcing in regulating retreat of the marine-based Irish-Sea Ice Stream*. *Geological Society of America Bulletin* DOI: 10.1130/B31852.1
- Smith MJ, Knight J. 2011. Palaeoglaciology of the last Irish ice sheet reconstructed from striae evidence. *Quaternary Science Reviews* **30**: 147-160.
- Smedley RK, Chiverrell RC, Ballantyne CK *et al.* 2017. Internal dynamics condition centennial-scale oscillations in marine-based ice-stream retreat. *Geology* **45**: 787–790.
- Stephen N, Synge FM. 1965. Late-Pleistocene shorelines and drift limits in north Donegal. *Proceedings of the Royal Irish Academy* **64B**: 131-153.
- Stone JO. 2000. Air pressure and cosmogenic isotope production. *Journal of Geophysical Research: Solid Earth* **105**: 23753-23759.
- Stone JO, Allan GL, Fifield KL *et al.* 1996. Cosmogenic chlorine-36 from calcium spallation. *Geochimica et Cosmochimica Acta* **60**: 679-692.
- Van Asch N, Lutz AF, Duijkers MCH *et al.* 2012. Rapid climate change during the Weichselian Lateglacial in Ireland: Chironomid-inferred summer temperatures from Fiddaun, C. Galway. *Palaeogeography, Palaeoclimatology, Palaeoecology* **315–316**: 1–11.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

769 Watson JE, Brooks SJ, Whitehouse NJ *et al.* 2010. Chironomid-inferred late-glacial summer air
770 temperatures from Lough Nadourcan, Co. Donegal, Ireland. *Journal of Quaternary Science* **25**:
771 1200-1210.
772
773 Wilcken KM, Freeman SPHT, Schnabel C *et al.* 2013. ³⁶Cl accelerator mass spectrometry with a
774 bespoke instrument. *Nuclear Instruments and Methods in Physics Research B* **294**: 107-114
775
776 Xu S, Dougans AB, Freeman SPHT *et al.* 2010. Improved ¹⁰Be and ²⁶Al AMS with a 5 MV
777 spectrometer. *Nuclear Instruments and Methods in Physics Research B* **268**: 736-738.

Figures:

Figure 1. Location map of Donegal and parts of Sligo and Mayo with terrestrial legacy ages (TCN and ^{14}C), BRITICE-CHRONO ^{14}C ages for the Donegal Bay Moraine, with core numbers, and sites of new TCN ages reported in this paper. Only those legacy ages of relevance to deglaciation are shown. The first TCN age for each site was calculated using the LLPR, the age in brackets was calculated using CRONUScalc. Ages are mean values of two or more ages for each site except for Glencolumbkille, which is represented, by a single age. Terrestrial ^{14}C ages are minimum ages for deglaciation. Ice divides and generalised ice flow directions are from Greenwood and Clark (2009a). Inset shows location of Donegal and the LGM limit of the BIIS.

Figure 2. **a:** Some of the granite boulders of the moraine at Bloody Foreland; **b:** Part of the spread of glacially-transported granite boulders at Rosguill; **c:** Rosguill boulder ROS-01. The survey pole is divided at intervals of 0.2 m; **d:** Sampled bedrock at Malin Head, site MH-02; **e:** Glencolumbkille boulder GC-02 undergoing sampling of projecting quartz vein; **f:** Blue Stack Mountains boulder BS-01, showing projecting quartz pebbles. Scale bar is divided at intervals of 0.06 m.

Figure 3. Location map of Donegal and parts of Sligo and Mayo with all ages (legacy and new) of relevance to deglaciation. The new TCN ages are given in the same format as the legacy ages in Figure 1. Note that the Malin Head age is a single age and the Glencolumbkille age is the mean of three new ages and the legacy age given in Figure 1. Inferred ice margins (isochrones) are shown as solid lines for ~26-25 ka (ILGM), 22 ka, ~21 ka, ~18 ka, ~16 ka and ~15 ka, and as broken lines for readvance limits at ~19.5 ka (Donegal Bay Moraine) and ~18 ka (Corvish).

Figure 4. Equal-area Gaussian probability distributions representing the uncertainty-weighted means and associated uncertainties for the cosmogenic ^{10}Be exposure ages obtained for samples from the ‘coastal’ sites on Aran Island ($n = 2$) and Bloody Foreland ($n = 7$), and the ‘inland mountain’ sites of Errigal Col ($n = 3$) and Poisoned Glen ($n = 2$). These distributions illustrate the overlap in the ages obtained for the two ‘coastal’ sites and for the two ‘inland mountain’ sites, and also the temporal interval of ≥ 4 ka that separates the two sets of ages.

Tables:

Table 1. Terrestrial legacy ages and BRITICE-CHRONO ages pertaining to the deglaciation of Donegal, Donegal Bay and north Mayo.

Table 2. Details of samples for TCN dating from Donegal.

Table 3. TCN (^{10}Be and ^{36}Cl) data and surface exposure ages with total uncertainties at 1σ for the Donegal samples. Analytical uncertainties (1σ) are given in parentheses.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47

Table 1. Terrestrial legacy ages and BRITICE-CHRONO ages pertaining to the deglaciation of Donegal, Donegal Bay and north Mayo.

Site	¹⁴ C age±1σ (cal. ka BP ¹)	¹⁰ Be age±1σ (LLPR ²)	¹⁰ Be age±1σ (CRONUScal ³)	³⁶ Cl ages±1σ ⁴	Material and context	Reference
DONEGAL						
<i>Malin Head</i>				25.1±1.1	Glacially-smoothed quartzite bedrock	Bowen <i>et al.</i> (2002)
<i>Bloody Foreland</i>				31.0±17.0	Not specified, but granite bedrock or boulder	Bowen <i>et al.</i> (2002)
<i>Corvish</i>	17.06±0.18 19.03±0.19 18.32±0.18 19.50±0.50 18.24±0.13 20.68±0.16				Marine microfauna: <i>Elphidium clavatum</i> Marine microfauna: <i>Elphidium clavatum</i> Marine microfauna: <i>Elphidium clavatum</i> Marine microfauna: <i>Elphidium clavatum</i> Marine microfauna: <i>Elphidium clavatum</i> Marine microfauna: <i>Elphidium clavatum</i>	McCabe and Clark (2003)
<i>Bloody Foreland</i>		21.4±1.4 (1.0) 18.6±1.2 (0.8)	20.7±1.9 (1.0) 18.0±1.6 (0.8)		Glacially-transported granite boulder Glacially-transported granite boulder	Ballantyne <i>et al.</i> (2007)
<i>Bloody Foreland</i>		18.0±1.8 (1.6) 34.1±3.1 (2.7) 22.0±1.8 (1.5) 21.4±1.9 (1.7) 21.4±2.1 (1.9) 23.8±2.2 (1.9) 21.9±2.3 (2.0) 22.3±2.3 (2.0)	17.5±2.2 (1.7) 32.5±3.7 (2.7) 21.3±2.3 (1.6) 20.7±2.3 (1.7) 20.7±2.5 (1.9) 23.0±2.6 (2.0) 21.2±2.7 (2.1) 21.6±2.7 (2.1)		Glacially-transported granite boulder Glacially-transported granite boulder Glacially-transported granite boulder Glacially-transported granite boulder Glacially-transported granite boulder Glacially-transported granite boulder Glacially-transported granite boulder Glacially-transported granite boulder	Clark <i>et al.</i> (2009a)
	<i>Mean</i> ^{5, 6}	21.8±1.2	21.1±1.8			
<i>Aran Island</i>		21.8±1.2 (0.7) 21.6±1.2 (0.7)	21.1±1.8 (0.8) 20.9±1.8 (0.8)		Glacially-transported granite boulder Granite bedrock	Cullen (2013)
	<i>Mean</i> ⁶	21.7±1.1	21.0±1.7			
<i>Glencolumbkille</i> ⁷		17.9±0.9 (0.5) 19.8±1.0 (0.5)	17.4±1.4 (0.5) 19.1±1.5 (0.6)		Vein quartz in glacially-transported schist boulder Vein quartz in schist roche moutonnée	Ballantyne <i>et al.</i> (2007)

Table 1 continued.

Errigal col	17.4±1.0 (0.6)	17.2±1.5 (0.6)	Glacially-plucked quartzite bedrock	Ballantyne <i>et al.</i> (2013)
	18.0±1.0 (0.6)	17.8±1.5 (0.6)	Glacially-plucked quartzite bedrock	
	17.9±1.0 (0.6)	17.7±1.5 (0.6)	Glacially-plucked quartzite bedrock	
Mean⁶	17.8±0.9	17.6±1.4		
Slieve League	16.9±1.0 (0.7)	16.7±1.4 (0.7)	Quartzite boulder from rockslope-failure debris	Ballantyne <i>et al.</i> (2013)
	17.6±1.2 (0.9)	17.4±1.6 (0.9)	Quartzite boulder from rockslope-failure debris	
	16.9±1.1 (0.9)	16.7±1.5 (0.8)	Quartzite boulder from rockslope-failure debris	
Mean⁶	17.1±0.9	16.9±1.4		
Lough Nadourcan	15.38±0.12		Organic mud, basal lake sediment, bulk sample	Watson <i>et al.</i> (2010)
NORTH MAYO				
Fiddauntawnanoneen	20.38±0.31		Marine shell: <i>Macoma calcarea</i>	McCabe <i>et al.</i> (1986)
Belderg Pier	19.16±0.21		Marine shell: <i>Macoma calcarea</i>	McCabe <i>et al.</i> (1986, 2005)
	19.23±0.26		Marine shell: <i>Macoma calcarea</i>	
	19.51±0.29		Marine microfauna: <i>Elphidium clavatum</i>	
	19.77±0.35		Marine microfauna: <i>Elphidium clavatum</i>	
	19.88±0.33		Marine shell: <i>Macoma calcarea</i>	
	19.92±0.34		Marine shell: <i>Macoma calcarea</i>	
	22.09±0.28		Marine microfauna: <i>Quinqueloculina seminulum</i>	
Donegal Bay	20.24±0.24		Mixed benthic foraminifera	Ó Cofaigh <i>et al.</i> (2018)
	17.92±0.16		Mixed benthic foraminifera	
Ox Mountains	17.0±1.6 (1.4)	17.0±2.0 (1.5)	Vein quartz in glacially-transported gneissic boulder	Clark <i>et al.</i> (2009b)
	15.7±1.6 (1.4)	16.0±2.0 (1.6)	Vein quartz in glacially-transported gneissic boulder	
	16.4±1.5 (1.3)	16.5±1.9 (1.4)	Vein quartz in glacially-transported gneissic boulder	
	16.9±1.5 (1.3)	17.0±1.9 (1.4)	Vein quartz in glacially-transported gneissic boulder	
	17.0±1.8 (1.7)	17.1±2.3 (1.8)	Vein quartz in glacially-transported gneissic boulder	
Mean⁶	16.6±1.0	16.7±1.5		
	19.1±1.8 (1.6)	19.2±2.3 (1.7)	Vein quartz in glacially-transported gneissic boulder	Clark <i>et al.</i> (2009b)
	21.1±1.7 (1.4)	21.0±2.3 (1.6)	Vein quartz in glacially-transported gneissic boulder	
Table 1 continued.				
	20.6±2.1 (1.8)	20.7±2.5 (2.0)	Vein quartz in glacially-transported gneissic boulder	

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47

826

827
828

<i>Mean</i> ⁶	<i>20.3±1.3</i>	<i>20.3±1.9</i>
--------------------------	-----------------	-----------------

¹Ages calculated using OxCal 4.2; Marine-13 dataset used for Corvish and Donegal Bay samples (Bronk Ramsey, 2009; Reimer *et al.*, 2013).
²Ages calculated using the Loch Lomond Production Rate, Lm scaling and 1 mm ka⁻¹ surface erosion rate. Uncertainties are total (external) uncertainties. Analytical uncertainties are given in parentheses.
³Ages calculated using CRONUScalc, Lm scaling and 1 mm ka⁻¹ surface erosion rate. Uncertainties are total (external) uncertainties. Analytical uncertainties are given in parentheses.
⁴Chlorine-36 ages cited as published by original authors.
⁵Mean derives from six consistent values reported by Clark *et al.* (2009a) and one age reported by Ballantyne *et al.* (2007).
⁶Uncertainty-weighted mean.
⁷Site is called Malin Beg in Ballantyne *et al.* (2007).

Table 2. Details of samples for TCN dating from Donegal.

Sample code	Grid reference	Latitude (°N)	Longitude (°W)	Altitude (m OD)	Thickness (cm)	Density (g cm ⁻³)	Topographic shielding	Material and context
North Donegal								
<i>Rosguill</i>								
ROS-01	C 0999 4222	55.22690	7.84304	65	4.0	2.66	0.9938	Glacially-transported granite boulder
ROS-02	C 1014 4203	55.22520	7.84062	105	5.0	2.67	0.9997	Glacially-transported granite boulder
ROS-04	C 1015 4191	55.22412	7.84055	105	3.0	2.66	0.9967	Glacially-transported granite boulder
<i>Malin Head</i>								
MH-02	C 3977 5955	55.38112	7.37255	65	5.0	2.59	0.9939	Ice-scoured quartzite bedrock
MH-03	C 3947 5960	55.38156	7.37716	30	3.0	2.65	0.9996	Vein quartz in quartzite bedrock
MH-04	C 3964 5946	55.38033	7.37458	55	2.5	2.58	0.6380	Ice-scoured quartzite bedrock
<i>Poisoned Glen</i>								
PG-01	B 9317 1863	55.01505	8.10675	73	2.0	2.61	0.9962	Glacially-transported granite boulder
PG-04	B 9319 1862	55.01495	8.10653	73	3.0	2.62	0.9891	Glacially-transported granite boulder
PG-05	B 9324 1862	55.01498	8.10572	75	2.5	2.56	0.9969	Glacially-transported granite boulder
South Donegal								
<i>Glencolumbkille</i>								
GCS-01	G 5100 8474	54.70850	8.76100	25	1.5	2.65	0.9974	Vein quartz in glacially-transported schist boulder
GCS-02	G 5107 8468	54.70790	8.75990	40	2.0	2.65	0.9983	Vein quartz in glacially-transported schist boulder
GCS-03	G 5114 8464	54.70760	8.75890	35	1.0	2.65	0.9982	Vein quartz in glacially-transported schist boulder

1										
2										
3	831	Table 2 continued.								
4	832									
5	833									
6	834									
7	835	Kilcar								
8	836	KC-01	G 6063 7468	54.61870	8.60960	50	5.0	3.05	0.9980	Glacially-transported dolerite boulder
9	837	KC-02	G 6063 7468	54.61870	8.60960	50	4.0	3.04	0.9987	Glacially-transported dolerite boulder
10	838	KC-03	G 6063 7468	54.61870	8.60960	50	3.5	3.04	0.9665	Glacially-transported dolerite boulder
11	839	KC-04	G 6066 7470	54.61890	8.60900	45	6.0	3.03	0.9838	Glacially-transported dolerite boulder
12	840									
13	841	Blue Stacks								
14	842	BS-01	G 9276 8608	54.72280	8.11320	150	1.0	2.65	0.9988	Quartz pebbles in glacially-transported conglomerate boulder
15	843	BS-02	G 9270 8604	54.72250	8.11400	148	3.5	2.40	0.9967	Glacially-transported conglomerate sandstone boulder
16	844	BS-03	G 9254 8611	54.72310	8.11650	150	2.0	2.65	0.9993	Quartz pebbles in glacially-transported conglomerate boulder
17	845	BS-04	G 9242 8620	54.72390	8.11840	163	5.0	2.26	0.9993	Glacially-transported conglomerate sandstone boulder
18	846									
19	847									
20	848									
21	849									
22	850									
23										
24										
25										
26										
27										
28										
29										
30										
31										
32										
33										
34										
35										
36										
37										
38										
39										
40										
41										
42										
43										
44										
45										
46										
47										

Table 3. Cosmogenic (^{10}Be and ^{36}Cl) data and surface exposure ages with total uncertainties at 1σ for the Donegal samples.Analytical uncertainties (1σ) are given in parentheses.

Sample code	AMS ID	^{10}Be (10^4 atoms g^{-1})	Blank ^{10}Be (10^4 atoms)	^{36}Cl (10^4 atoms g^{-1})	Blank ^{36}Cl (10^4 atoms)	^{10}Be Exposure Age ¹ (LLPR)	^{10}Be Exposure Age ² (CRONUScale)	^{36}Cl Exposure Age ³ (CRONUScale)
North Donegal								
<i>Rosguill</i>								
ROS-01	b10320	7.828±0.253	0.609±0.102			18.7±1.1 (0.7)	18.4±1.6 (0.7)	
ROS-02	b10322	9.257±0.418	0.609±0.102			21.4±1.4 (1.0)	21.0±1.9 (1.0)	
ROS-04	b10323	8.331±0.254	0.609±0.102			18.9±1.1 (0.6)	18.7±1.6 (0.6)	
	<i>Mean</i> ^{4, 5}					18.8±0.9	18.6±1.5	
<i>Malin Head</i>								
MH-02	b10324	9.602±0.363	0.609±0.102			23.3±1.4 (1.0)	23.0±2.0 (0.9)	
MH-03	b10286	8.433±0.272	0.609±0.102			20.7±1.2 (0.7)	20.2±1.7 (0.7)	
MH-04	b10426	6.881±0.472	0.827±0.142			25.5±1.7 (1.2)	25.2±2.3 (1.2)	
<i>Poisoned Glen</i>								
PG-01	b10628	7.413±0.328	0.785±0.108			17.2±1.1 (0.8)	16.7±1.5 (0.7)	
PG-04	b10629	6.876±0.315	0.785±0.108			16.2±1.0 (0.8)	15.7±1.5 (0.7)	
PG-05	b10630	5.605±0.276	0.785±0.108			13.0±0.9 (0.6)	12.6±1.2 (0.6)	
	<i>Mean</i> ^{4, 6}					16.7±0.9	16.2±1.4	
South Donegal								
<i>Glencolumbkille</i>								
GCS-02	b8570	6.672±0.291	0.894±0.159			16.3±1.0 (0.7)	16.0±1.4 (0.7)	
GCS-03	b8572	7.192±0.306	0.894±0.159			17.3±1.1 (0.7)	17.0±1.5 (0.7)	
GCS-04	b8573	6.873±0.293	0.894±0.159			16.5±1.0 (0.7)	16.2±1.5 (0.7)	

Table 3 continued.

1						
2						
3						
4		<i>Mean</i> ⁴			<i>16.7±0.8</i>	<i>16.4±1.4</i>
5		<i>Mean</i> ^{4, 7}			<i>17.2±0.8</i>	<i>16.8±1.3</i>
6						
7						
8	<i>Kilcar</i>					
9	KC-01	c4058		11.767±0.451	6.872±1.722	18.1±1.7 (0.7)
10	KC-02	c4059		12.378±0.598	6.872±1.722	37.2±6.2 (2.1)
11	KC-03	c4060		11.796±0.473	6.872±1.722	42.0±6.0 (1.9)
12	KC-04	c4061		13.951±0.558	6.872±1.722	37.1±5.3 (1.7)
13						
14	<i>Blue Stacks</i>					
15						
16	BS-01	b9962	6.163±0.331	1.683±0.177	14.6±0.9 (0.6)	14.5±1.3 (0.6)
17	BS-02	b8568	7.097±0.301	0.894±0.159	15.6±0.9 (0.7)	15.6±1.4 (0.6)
18	BS-03	b8569	6.984±0.303	0.894±0.159	14.9±0.9 (0.6)	14.8±1.3 (0.6)
19	BS-04	b10285	6.729±0.206	0.814±0.103	14.4±0.8 (0.5)	14.5±1.3 (0.5)
20		<i>Mean</i> ⁴			<i>14.8±0.7</i>	<i>14.8±1.2</i>
21						
22						

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

851

¹Exposure age based on Loch Lomond Production Rate (LLPR), time-dependent Lm scaling (Lal, 1991; Stone, 2000) and assuming 1 mm ka⁻¹ erosion.

²Exposure age derived from CRONUScal v2.0 with Lm scaling and assuming 1 mm ka⁻¹ erosion.

³Exposure age based on CRONUScal v2.0 with Lm scaling.

⁴Uncertainty-weighted mean value.

⁵Mean value based on ROS-01 and -04 only.

⁶Mean value based on PG-01 and -04 only.

⁷Mean value includes an additional sample age reported by Ballantyne *et al.* (2007).



Figure 2

338x190mm (96 x 96 DPI)




Figure 3




Figure 4

136x92mm (300 x 300 DPI)